

IR spectroscopy of pre-cataclysmic binaries

C. Tappert¹, B. T. Gänsicke², R. E. Mennickent³

¹*Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica, Casilla 306, Santiago 22, Chile*

²*Department of Physics, University of Warwick, Coventry CV4 7AL, UK*

³*Grupo de Astronomía, Universidad de Concepción, Casilla 160-C, Concepción, Chile*

Abstract. We have taken near-infrared (*JHK*) spectroscopic data of a sample of pre-CVs, in order to examine them for signatures of nuclear evolution. Here we present preliminary results for the three systems BPM 71214, RR Cae, and RE 1016–053.

The data were obtained with SofI at the NTT, La Silla. The low resolution Grism Blue for *J* ($R \sim 800$) and medium resolution grisms for *H* ($R \sim 1200$) and *K* ($R \sim 1300$) were used. The standard reduction process included correction for telluric absorption, but no flux calibration. Equivalent widths were measured using the wavelength ranges given by Wallace et al. (2000), Meyer et al. (1998), and Wallace & Hinkle (1997). For the *J* spectra the continuum was determined using the regions from Wallace et al. (2000), with the exception of CN. For this line and for the *H* and *K* data, the equivalent widths were measured with respect to linear fits to the local continuum.

The results are plotted in Fig. 1 together with standard stars of luminosity classes V and III. We find that within the errors there are no systematic deviations from the values for main-sequence stars, although a few differences for individual lines do exist. However, a proper comparison is somewhat problematic, due to the low number of standard stars with spectral types later than M0, and due to the low efficiency of the SofI *K* grism longward of $2.3\mu\text{m}$ ($<4350\text{ cm}^{-1}$), which practically inhibits the measurement of those CO bands that have been found to show systematic anomalous abundances in CVs (Harrison et al. 2004). We therefore plan to supplement our survey with *K* spectra from ISAAC at the VLT.

References

- Harrison, T.E., Osborne, H.L., Howell, S.B., 2004, AJ, 127, 3493
- Meyer, M.R., Edwards, S., Hinkle, K.H., Strom, S.E., 1998, ApJ, 508, 397
- Wallace, L., Hinkle, K., 1997, ApJS, 111, 445
- Wallace, L., Meyer, M.R., Hinkle, K., Edwards, S., 2000, ApJ, 535, 325




Figure 1. Equivalent widths in cm^{-1} of selected regions in the JHK spectra. Standard star data for J and H have been taken from Wallace et al. (2000) and Meyer et al. (1998), respectively. Equivalent widths for K standards have been measured using electronically available spectra from Wallace & Hinkle (1997). In the plots, the small symbols indicate the standard data: dots for dwarfs and crosses for giants. The pre-CVs are symbolized as follows: Square = RE 1016-053, Triangle = BPM 71214, Diamond = RR Cae. Horizontal error bars correspond to the uncertainty of the spectral type given in the literature, vertical error bars give the standard deviation of the average equivalent widths measured from five individual fits to the local continuum.